

# Topic 1 // Lesson 04

{ To determine an election using *Pairwise Comparisons*

& Like a Round Robin Tournament in which every player plays once against every other, where *head-to-head* matches are called **pairwise comparisons**.

Method of Pairwise Comparisons

- ⌘· Each candidate goes head-to-head and winner gets a single point
- ⌘· If a tie, then both get 1/2 point
- ⌘· Winner is candidate with most points
- ⌘· Ties are very common in this method

## Method of Pairwise Comparisons

|             |    |    |   |   |   |
|-------------|----|----|---|---|---|
| # of voters | 14 | 10 | 8 | 4 | 1 |
| 1st         | A  | C  | D | B | C |
| 2nd         | B  | B  | C | D | D |
| 3rd         | C  | D  | B | C | B |
| 4th         | D  | A  | A | A | A |

A vs B:

A vs C:

A vs D:

B vs C:

B vs D:

C vs D:

Overall Tally: A =    B =    C =    D =

The Winner is:

# Method of Pairwise Comparisons

|             |    |    |   |   |   |
|-------------|----|----|---|---|---|
| # of voters | 14 | 10 | 8 | 4 | 1 |
| 1st         | A  | C  | D | B | C |
| 2nd         | B  | B  | C | D | D |
| 3rd         | C  | D  | B | C | B |
| 4th         | D  | A  | A | A | A |

A vs B: 14 to 23, B=1

A vs C: 14 to 23, C=1

A vs D: 14 to 23, D=1

B vs C: 18 to 19, C=1

B vs D: 28 to 9, B=1

C vs D: 25 to 12, C=1

Overall Tally: A = 0 B= 2 C= 3 D= 1

The Winner is: **Person C**

# Method of Pairwise Comparisons

|             |   |   |   |   |   |   |   |
|-------------|---|---|---|---|---|---|---|
| # of voters | 2 | 6 | 4 | 1 | 1 | 4 | 4 |
| 1st         | A | B | B | C | C | D | E |
| 2nd         | D | A | A | B | D | A | C |
| 3rd         | C | C | D | A | A | E | D |
| 4th         | B | D | E | D | B | C | B |
| 5th         | E | E | C | E | E | B | A |

Overall Tally: A =    B =    C =    D =

The Winner is:

Ex 1.12 LAXer's Draft Choice Election

|             |   |   |   |   |   |   |   |
|-------------|---|---|---|---|---|---|---|
| # of voters | 2 | 6 | 4 | 1 | 1 | 4 | 4 |
| 1st         | A | B | B | B | D | D | E |
| 2nd         | D | A | A | A | A | A | D |
| 3rd         | B | D | D | D | B | E | B |
| 4th         | E | E | E | E | E | B | A |

Overall Tally: A =    B =    C =    D =  
The Winner is:

Ex 1.12 LAXer's Draft Choice Election

If Candidate X is a winner of an election and in a recount one of the *nonwinning* candidates withdraws or is disqualified, then X should still be a winner of the election.

Alternative Interpretation: If Candidate X is a winner of an election and in a reelection another candidate that has no chance of winning (an “irrelevant alternative”) enters the race, then X should still be the winner

## **Independence of Irrelevant Alternatives Criterion (IIA)**



Add up 1 to 99.

**How many Pairwise Comparisons?**

Gauss

**How many Pairwise Comparisons?**

# Sum of Consecutive Integers Formula

$$1 + 2 + 3 + \dots + L = \frac{L(L + 1)}{2}$$

Consider election with 10 candidates:

A, B, C, D, E, F, G, H, I, and J

Compare A with 9 others is 9 comparisons

Compare B with 8 others is 8 comparisons

Compare C with 7 others is 7 comparisons

Compare D with 6 others is 6 comparisons

Continuing until J...

Total #:  $1 + 2 + 3 + \dots + 8 + 9 =$

## Counting Pairwise Comparisons

# Number of Pairwise Comparisons

In an election with  $N$  candidates the total number of pairwise comparisons is

$$\frac{(N - 1)N}{2}$$

Practice 1. How many comparisons would occur in a 12 candidate race?  
Practice 2. How many comparisons would occur in a 20 candidate race?